



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 35/76	A1	(11) International Publication Number: WO 99/08692 (43) International Publication Date: 25 February 1999 (25.02.99)
(21) International Application Number: PCT/CA98/00774 (22) International Filing Date: 12 August 1998 (12.08.98) (30) Priority Data: 08/911,383 13 August 1997 (13.08.97) US (71)(72) Applicants and Inventors: LEE, Patrick, W., K. [CA/CA]; 22 Varmoor Place, N.W. Calgary, Alberta T3A 0A1 (CA). STRONG, James [CA/CA]; 5071 Nesbitt Road, N.W. Calgary, Alberta T2K 2N6 (CA). COFFEY, Mathew, C. [CA/CA]; 536 34A Street, N.W. Calgary, Alberta T2N 2Y5 (CA). (74) Agents: VAN ZANT, Joan, M. et al.; Van Zant & Associates, Suite 1407, 77 Bloor Street West, Toronto, Ontario M5S 1M2 (CA).		(81) Designated States: AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CU, CZ, DE, DK, EE, ES, FI, GB, GE, GH, GM, HR, HU, ID, IL, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, UA, UG, US, UZ, VN, YU, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i>
(54) Title: REOVIRUS FOR THE TREATMENT OF NEOPLASIA (57) Abstract Methods for treating neoplasia, by administering reovirus to a Ras-mediated neoplasm, and use of reovirus for manufacture of a medicament for the treatment of neoplasia, are disclosed. The reovirus is administered so that it ultimately directly contacts cells of the neoplasm. Human reovirus, non-human mammalian reovirus, and/or avian reovirus can be used. If the reovirus is human reovirus, type 1 (e.g., strain Lang), type 2 (e.g., strain Jones), type 3 (e.g., strain Dearing or strain Abney), as well as other serotypes or strains of reovirus can be used. Combinations of more than one type and/or strain of reovirus can be used, as can reovirus from different species of animal. Either solid neoplasms or hematopoietic neoplasms can be treated.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

REOVIRUS FOR THE TREATMENT OF NEOPLASIA

RELATED APPLICATION

This application is a Continuation-in-Part of U.S. Serial No. 08/911,383 filed on August 13, 1997, the entire teachings of which are incorporated herein by reference.

5 BACKGROUND OF THE INVENTION

Normal cell proliferation is regulated by a balance between growth-promoting proto-oncogenes and growth-constraining tumor-suppressor genes. Tumorigenesis can be caused by genetic alterations to the genome that result in the mutation of those cellular elements that govern the interpretation of cellular signals, such as potentiation
10 of proto-oncogene activity or inactivation of tumor suppression. It is believed that the interpretation of these signals ultimately influences the growth and differentiation of a cell, and that misinterpretation of these signals can result in neoplastic growth (neoplasia).

Genetic alteration of the proto-oncogene Ras is believed to contribute to
15 approximately 30% of all human tumors (Wiessmuller, L. and Wittinghofer, F. (1994), *Cellular Signaling* 6(3):247-267; Barbacid, M. (1987) *A Rev. Biochem.* 56, 779-827). The role that Ras plays in the pathogenesis of human tumors is specific to the type of tumor. Activating mutations in Ras itself are found in
most types of human malignancies, and are highly represented in pancreatic cancer
20 (80%), sporadic colorectal carcinomas (40-50%), human lung adenocarcinomas (15-24%), thyroid tumors (50%) and myeloid leukemia (30%) (Millis, NE *et al.* (1995)

-2-

Cancer Res. 55:1444; Chaubert, P. *et al.* (1994), *Am. J. Path.* 144:767; Bos, J. (1989) *Cancer Res.* 49:4682). Ras activation is also demonstrated by upstream mitogenic signaling elements, notably by tyrosine receptor kinases (RTKs). These upstream elements, if amplified or overexpressed, ultimately result in elevated Ras activity by the
5 signal transduction activity of Ras. Examples of this include overexpression of PDGFR in certain forms of glioblastomas, as well as in c-erbB-2/neu in breast cancer (Levitzki, A. (1994) *Eur. J. Biochem.* 226:1; James, P.W., *et al.* (1994) *Oncogene* 9:3601; Bos, J. (1989) *Cancer Res.* 49:4682).

Current methods of treatment for neoplasia include surgery, chemotherapy and
10 radiation. Surgery is typically used as the primary treatment for early stages of cancer; however, many tumors cannot be completely removed by surgical means. In addition, metastatic growth of neoplasms may prevent complete cure of cancer by surgery. Chemotherapy involves administration of compounds having antitumor activity, such as alkylating agents, antimetabolites, and antitumor antibiotics. The efficacy of
15 chemotherapy is often limited by severe side effects, including nausea and vomiting, bone marrow depression, renal damage, and central nervous system depression. Radiation therapy relies on the greater ability of normal cells, in contrast with neoplastic cells, to repair themselves after treatment with radiation. Radiotherapy cannot be used to treat many neoplasms, however, because of the sensitivity of tissue surrounding the
20 tumor. In addition, certain tumors have demonstrated resistance to radiotherapy and such may be dependent on oncogene or anti-oncogene status of the cell (Lee, J.M. *et al.* (1993) *PNAS* 90:5742-5746; Lowe, S.W. *et al.* (1994) *Science*, 266:807-810; Raybaud-Diogene, H. *et al.* (1997) *J. Clin. Oncology*, 15(3):1030-1038). In view of the drawbacks associated with the current means for treating neoplastic growth, the need
25 still exists for improved methods for the treatment of most types of cancers.

-3-

SUMMARY OF THE INVENTION

The present invention pertains to methods for treating neoplasia in a mammal, using reovirus, and to use of reovirus for manufacture of a medicament for the treatment of neoplasia. Reovirus is administered to a neoplasm, in which an element of the Ras signaling pathway (either upstream or downstream) is activated to an extent that results in reovirus-mediated oncolysis of cells of the neoplasm. The reovirus can be administered in a single dose or in multiple doses; furthermore, more than one neoplasm in an individual mammal can be treated concurrently. Both solid neoplasms and hematopoietic neoplasms can be targeted. The reovirus is administered so that it contacts cells of the mammal (e.g., by injection directly into a solid neoplasm, or intravenously into the mammal for a hematopoietic neoplasm). The methods can be used to treat neoplasia in a variety of mammals, including mice, dogs, cats, sheep, goats, cows, horses, pigs, and non-human primates. Preferably, the methods are used to treat neoplasia in humans.

The methods and uses of the invention provide an effective means to treat neoplasia, without the side effects associated with other forms of cancer therapy. Furthermore, because reovirus is not known to be associated with disease, any safety concerns associated with deliberate administration of a virus are minimized.

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is a depiction of the molecular basis of reovirus oncolysis, in which the reovirus usurps the host cell Ras signalling pathway.

Figure 2 is a graphic representation of the effects over time of active (open circles) or inactivated (closed circles) reovirus serotype 3 (strain Dearing) on the size of murine THC-11 tumors grown in severe combined immunodeficiency (SCID) mice.

-4-

The plotted values represent the mean of the measurements with the standard error of the mean also shown.

Figure 3 is a graphic representation of the effects over time of active (open circles) or inactivated (closed circles) reovirus serotype 3 (strain Dearing) on the size of human glioblastoma U-87 xenografts grown in SCID mice. The plotted values represent the mean of the measurements with the standard error of the mean also shown.

Figure 4 is a graphic representation of the effects over time of active (open circles, open squares) or inactivated (closed circles, closed squares) reovirus serotype 3 (strain Dearing) on the size of treated (circles) or untreated (squares) bilateral human glioblastoma U-87 xenografts grown in SCID mice. The plotted values represent the mean of the measurements with the standard error of the mean also shown.

The foregoing and other objects, features and advantages of the invention will be apparent from the following more particular description of preferred embodiments of the invention, as illustrated in the accompanying drawings in which like reference characters refer to the same parts throughout the different views. The drawings are not necessarily to scale, emphasis instead being placed upon illustrating the principles of the invention.

DETAILED DESCRIPTION OF THE INVENTION

The invention pertains to methods of treating a neoplasm in a mammal, by administering reovirus to the neoplasm. The name reovirus (*Respiratory and enteric orphan virus*) is a descriptive acronym suggesting that these viruses, although not associated with any known disease state in humans, can be isolated from both the respiratory and enteric tracts (Sabin, A.B. (1959), *Science* 130:966). The mammalian reovirus consists of three serotypes: type 1 (strain Lang or T1L), type 2 (strain Jones, T2J) and type 3 (strain Dearing or strain Abney, T3D). The three serotypes are easily

-5-

identifiable on the basis of neutralization and hemagglutinin-inhibition assays (Sabin, A.B. (1959), *Science* 130:966; Fields, B.N. *et al.* (1996), *Fundamental Virology*, 3rd Edition, Lippincott-Raven; Rosen, L. (1960) *Am. J. Hyg.* 71:242; Stanley, N.F. (1967) *Br. Med. Bull.* 23:150).

- 5 Although reovirus is not known to be associated with any particular disease (Tyler, K.L. and Fields, B.N., in *Fields Virology* (Fields, B.N., Knipe, D.M., and Howley, P.M. eds), Lippincott-Raven, Philadelphia, 1996, p. 1597), many people have been exposed to reovirus by the time they reach adulthood (i.e., fewer than 25% in children < 5 years old, to greater than 50% in those 20-30 years old (Jackson G.G. and
- 10 Muldoon R.L. (1973) *J. Infect. Dis.* 128:811; Stanley N.F. (1974) *In: Comparative Diagnosis of Viral Diseases*, edited by E. Kurstak and K. Kurstak, 385-421, Academic Press, New York)..

- For mammalian reoviruses, the cell surface recognition signal is sialic acid (Armstrong, G.D. *et al.* (1984), *virology* 138:37; Gentsch, J.R.K. and Pacitti, A.F.
- 15 (1985), *J. Virol.* 56:356; Paul R.W. *et al.* (1989) *Virology* 172:382-385). Due to the ubiquitous nature of sialic acid, reovirus binds efficiently to a multitude of cell lines and as such can potentially target many different tissues; however, there are significant differences in susceptibility to reovirus infection between cell lines.

- As described herein, Applicants have discovered that cells which are resistant to
- 20 reovirus infection became susceptible to reovirus infection when transformed with a gene in the Ras pathway. "Resistance" of cells to reovirus infection indicates that infection of the cells with the virus did not result in significant viral production or yield. Cells that are "susceptible" are those that demonstrate induction of cytopathic effects, viral protein synthesis, and/or virus production. Resistance to reovirus infection was
- 25 found to be at the level of gene translation, rather than at early transcription: while viral transcripts were produced, virus proteins were not expressed. Viral gene transcription

-6-

in resistant cells correlated with phosphorylation of an approximately 65 kDa cell protein, determined to be double-stranded RNA-activated protein kinase (PKR), that was not observed in transformed cells. Phosphorylation of PKR lead to inhibition of translation. When phosphorylation was suppressed by 2-aminopurine, a known
5 inhibitor of PKR, drastic enhancement of reovirus protein synthesis occurred in the untransformed cells. Furthermore, in a severe combined immunodeficiency (SCID) mouse model in which tumors were created on both the right and left hind flanks revealed that reovirus significantly reduced tumor size when injected directly into the right-side tumor; in addition, significant reduction in tumor size was also noted on the
10 left-side tumor which was not directly injected with reovirus, indicating that the oncolytic capacity of the reovirus was systemic as well as local.

These results indicated that reovirus uses the host cell's Ras pathway machinery to downregulate PKR and thus reproduce. Figure 1 depicts the usurpation of the host cell Ras signalling pathway by reovirus. As shown in Figure 1, for both untransformed
15 (reovirus-resistant) and EGFR-, Sos-, or ras-transformed (reovirus-susceptible) cells, virus binding, internalization, uncoating, and early transcription of viral genes all proceed normally. In the case of untransformed cells, secondary structures on the early viral transcripts inevitably trigger the phosphorylation of PKR, thereby activating it, leading to the phosphorylation of the translation initiation factor eIF-2 α , and hence the
20 inhibition of viral gene translation. In the case of EGFR-, Sos-, or ras-transformed cells, the PKR phosphorylation step is prevented or reversed by Ras or one of its downstream elements, thereby allowing viral gene translation to ensue. The action of Ras (or a downstream element) can be mimicked by the use of 2-aminopurine (2-AP), which promotes viral gene translation (and hence reovirus infection) in untransformed cells by
25 blocking PKR phosphorylation.

-7-

Based upon these discoveries, Applicants have developed methods for treating neoplasms in mammals. Representative mammals include mice, dogs, cats, sheep, goats, cows, horses, pigs, non-human primates, and humans. In a preferred embodiment, the mammal is a human.

5 In the methods of the invention, reovirus is administered to a neoplasm in the individual mammal. Representative types of human reovirus that can be used include type 1 (e.g., strain Lang or T1L); type 2 (e.g., strain Jones or T2J); and type 3 (e.g., strain Dearing or strain Abney, T3D or T3A); other strains of reovirus can also be used. In a preferred embodiment, the reovirus is strain Dearing. Alternatively, the reovirus
10 can be a non-human mammalian reovirus (e.g., non-human primate reovirus, such as baboon reovirus; equine; or canine reovirus), or a non-mammalian reovirus (e.g., avian reovirus). A combination of different serotypes and/or different strains of reovirus, such as reovirus from different species of animal, can be used. The reovirus is "naturally-occurring": that is, it can be isolated from a source in nature and has not been
15 intentionally modified by humans in the laboratory. For example, the reovirus can be from a "field source": that is, from a human patient. If desired, the reovirus can be chemically or biochemically pretreated (e.g., by treatment with a protease, such as chymotrypsin or trypsin) prior to administration to the neoplasm. Such pretreatment removes the outer coat of the virus and may thereby result in better infectivity of the
20 virus.

 The neoplasm can be a solid neoplasm (e.g., sarcoma or carcinoma), or a cancerous growth affecting the hematopoietic system (a "hematopoietic neoplasm"; e.g., lymphoma or leukemia). A neoplasm is an abnormal tissue growth, generally forming a distinct mass, that grows by cellular proliferation more rapidly than normal tissue
25 growth. Neoplasms show partial or total lack of structural organization and functional coordination with normal tissue. As used herein, a "neoplasm", also referred to as a

-8-

"tumor", is intended to encompass hematopoietic neoplasms as well as solid neoplasms. At least some of the cells of the neoplasm have a mutation in which the Ras gene (or an element of the Ras signaling pathway) is activated, either directly (e.g., by an activating mutation in Ras) or indirectly (e.g., by activation of an upstream element in the Ras pathway). Activation of an upstream element in the Ras pathway includes, for example, transformation with epidermal growth factor receptor (EGFR) or Sos. A neoplasm that results, at least in part, by the activation of Ras, an upstream element of Ras, or an element in the Ras signalling pathway is referred to herein as a "Ras-mediated neoplasm". One neoplasm that is particularly susceptible to treatment by the methods of the invention is pancreatic cancer, because of the prevalence of Ras-mediated neoplasms associated with pancreatic cancer. Other neoplasms that are particularly susceptible to treatment by the methods of the invention include breast cancer, brain cancer (e.g., glioblastoma), lung cancer, prostate cancer, colorectal cancer, thyroid cancer, renal cancer, adrenal cancer, liver cancer, and leukemia.

The reovirus is typically administered in a physiologically acceptable carrier or vehicle, such as phosphate-buffered saline, to the neoplasm. "Administration to a neoplasm" indicates that the reovirus is administered in a manner so that it contacts the cells of the neoplasm (also referred to herein as "neoplastic cells"). The route by which the reovirus is administered, as well as the formulation, carrier or vehicle, will depend on the location as well as the type of the neoplasm. A wide variety of administration routes can be employed. For example, for a solid neoplasm that is accessible, the reovirus can be administered by injection directly to the neoplasm. For a hematopoietic neoplasm, for example, the reovirus can be administered intravenously or intravascularly. For neoplasms that are not easily accessible within the body, such as metastases or brain tumors, the reovirus is administered in a manner such that it can be transported systemically through the body of the mammal and thereby reach the

-9-

neoplasm (e.g., intrathecally, intravenously or intramuscularly). Alternatively, the reovirus can be administered directly to a single solid neoplasm, where it then is carried systemically through the body to metastases. The reovirus can also be administered subcutaneously, intraperitoneally, topically (e.g., for melanoma), orally (e.g., for oral or esophageal neoplasm), rectally (e.g., for colorectal neoplasm), vaginally (e.g., for cervical or vaginal neoplasm), nasally or by inhalation spray (e.g., for lung neoplasm).

The reovirus is administered in an amount that is sufficient to treat the neoplasm (e.g., an "effective amount"). A neoplasm is "treated" when administration of reovirus to cells of the neoplasm effects oncolysis of the neoplastic cells, resulting in a reduction in size of the neoplasm, or in a complete elimination of the neoplasm. The reduction in size of the neoplasm, or elimination of the neoplasm, is generally caused by lysis of neoplastic cells ("oncolysis") by the reovirus. The effective amount will be determined on an individual basis and may be based, at least in part, on consideration of the type of reovirus; the individual's size, age, gender; and the size and other characteristics of the neoplasm. For example, for treatment of a human, approximately 10^3 to 10^{12} plaque forming units (PFU) of reovirus can be used, depending on the type, size and number of tumors present. The reovirus can be administered in a single dose, or multiple doses (i.e., more than one dose). The multiple doses can be administered concurrently, or consecutively (e.g., over a period of days or weeks). The reovirus can also be administered to more than one neoplasm in the same individual.

The invention is further illustrated by the following Exemplification.

EXEMPLIFICATION

MATERIALS AND METHODS

Cells and Virus

-10-

Parental NIH-3T3 cell lines along with NIH-3T3 cells transformed with a number of oncogenes were obtained from a variety of sources. Parental NIH-3T3 and NIH-3T3 cells transfected with the Harvey-ras (H-ras) and EJ-ras oncogenes were a generous gift of Dr. Douglas Faller (Boston University School of Medicine). NIH-3T3
5 cells along with their Sos-transformed counterparts (designated TNIH#5) were a generous gift of Dr. Michael Karin (University of California, San Diego). Dr. H.-J. Kung (Case Western Reserve University) kindly donated parental NIH-3T3 cells along with NIH-3T3 cells transfected with the v-erbB oncogene (designated THC-11). 2H1 cells, a derivative of the C3H 10T1/2 murine fibroblast line, containing the Harvey-ras
10 gene under the transcriptional control of the mouse metallothionein-I promoter were obtained from Dr. Nobumichi Hozumi (Mount Sinai Hospital Research Institute). These 2H1 cells are conditional ras transformant that express the H-ras oncogene in the presence of 50 μ M ZnSO₄. All cell lines were grown in Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum (FBS).

15 The NIH-3T3 tet-myc cells were obtained from Dr. R.N. Johnston (University of Calgary) and were grown in DMEM containing 10% heat-inactivated FBS and antibiotics in the presence or absence of 2 μ g/ml tetracycline (Helbing, C.C. *et al.*, *Cancer Res.* 57:1255-1258 (1997)). In the presence of tetracycline, expression of the human *c-myc* gene is repressed. Removal of tetracycline results in the elevation of
20 expression of *c-myc* by up to 100-fold in these cells, which also display a transformed phenotype.

The PKR ^{+/+} and PKR ^{0/0} mouse embryo fibroblasts (MEFs) were obtained from Dr. B.R.G. Williams (the Cleveland Clinic Foundation) and were grown in α -MEM containing fetal bovine serum and antibiotics as previously described (Yang, Y.L. *et al.*
25 *EMBO J.* 14:6095-6106 (1995); Der, S.D. *et al.*, *Proc. Natl. Acad. Sci. USA* 94:3279-3283 (1997)).

-11-

The Dearing strain of reovirus serotype 3 used in these studies was propagated in suspension cultures of L cells and purified according to Smith (Smith, R.E. *et al.*, (1969) *Virology*, 39:791-800) with the exception that β -mercaptoethanol (β -ME) was omitted from the extraction buffer. Reovirus labelled with [35 S]methionine was grown and purified as described by McRae and Joklik (McRae, M.A. and Joklik, W.K., (1978) *Virology*, 89:578-593). The particle/PFU ratio for purified reovirus was typically 100/1.

Immunofluorescent analysis of reovirus infection

For the immunofluorescent studies the NIH-3T3, TNIH#5, H-ras, EJ-ras, 2H1 (+/- ZnSO₄), and THC-11 cells were grown on coverslips, and infected with reovirus at a multiplicity of infection (MOI) of ~10 PFU cell or mock-infected by application of the carrier agent (phosphate-buffered saline, PBS) to the cells in an identical fashion as the administration of virus to the cells. At 48 hours postinfection, cells were fixed in an ethanol/acetic acid (20/1) mixture for 5 minutes, then rehydrated by sequential washes in 75%, 50% and 25% ethanol, followed by four washes with phosphate-buffered saline (PBS). The fixed and rehydrated cells were then exposed to the primary antibody (rabbit polyclonal anti-reovirus type 3 serum diluted 1/100 in PBS) [antiserum prepared by injection of rabbits with reovirus serotype 3 in Freund's complete adjuvant, and subsequent bleedings] for 2 hours at room temperature. Following three washes with PBS, the cells were exposed to the secondary antibody [goat anti-rabbit IgG (whole molecule)-fluorescein isothiocyanate conjugate (FITC) [Sigma ImmunoChemicals F-0382] diluted 1/100 in PBS containing 10% goat serum and 0.005% Evan's Blue] for 1 hour at room temperature. Finally, the fixed and treated cells were washed three more times with PBS and then once with double-distilled water, dried and mounted on slides in 90% glycerol containing 0.1% phenylenediamine, and viewed with a Zeiss Axiophot

-12-

microscope on which Carl Zeiss camera was mounted (the magnification for all pictures was 200X).

Detection of MAP Kinase (ERK) Activity

The PhosphoPlus p44/42 MAP kinase (Thr202/Tyr204) Antibody kit (New
5 England Biolabs) was used for the detection of MAP kinase in cell lysates according to
the manufacturer's instructions. Briefly, subconfluent monolayer cultures were lysed
with the recommended SDS-containing sample buffer, and subjected to SDS-PAGE,
followed by electroblotting onto nitrocellulose paper. The membrane was then probed
with the primary antibody (anti-total MAPK or anti-phospho-MAPK), followed by the
10 horseradish peroxidase (HRP)-conjugated secondary antibody as described in the
manufacturer's instruction manual.

Radiolabelling of reovirus-infected cells and preparation of lysates

Confluent monolayers of NIH-3T3, TNIH#5, H-ras, EJ-ras, 2H1 (+/- ZnSO₄),
and THC-11 cells were infected with reovirus (MOI ~10 PFU/cell). At 12 hours
15 postinfection, the media was replaced with methionine-free DMEM containing 10%
dialyzed FBS and 0.1 mCi/ml [³⁵S]methionine. After further incubation for 36 hours at
37°C, the cells were washed in phosphate-buffered saline (PBS) and lysed in the same
buffer containing 1% Triton X-100, 0.5% sodium deoxycholate and 1 mM EDTA. The
nuclei were then removed by low speed centrifugation and the supernatants were stored
20 at -70°C until use.

Preparation of cytoplasmic extracts for in vitro kinase assays

Confluent monolayers of the various cell lines were grown on 96 well cell
culture plates. At the appropriate time postinfection the media was aspirated off and the

-13-

cells were lysed with a buffer containing 20mM HEPES [pH 7.4], 120 mM KCl, 5 mM MgCl₂, 1 mM dithiothreitol, 0.5% Nonidet P-40, 2 µg/ml leupeptin, and 50 µg/ml aprotinin. The nuclei were then removed by low-speed centrifugation and the supernatants were stored at -70°C until use.

- 5 Cytoplasmic extracts were normalized for protein concentrations before use by the Bio-Rad protein microassay method. Each in vitro kinase reaction contained 20 µl of cell extract, 7.5 µl of reaction buffer (20 mM HEPES [pH 7.4], 120 mM KCl, 5 mM MgCl₂, 1 mM dithiothreitol, and 10% glycerol) and 7.0 µl of ATP mixture (1.0 µCi[γ-³²P]ATP in 7 µl of reaction buffer), and was incubated for 30 minutes at 37°C
- 10 (Mundschau, L.J., and Faller, D.V., *J. Biol. Chem.*, 267:23092-23098 (1992)).
- Immediately after incubation the labelled extracts were either boiled in Laemmli SDS-sample buffer or were either precipitated with agarose-poly(I)poly(C) beads or immunoprecipitated with an anti-PKR antibody.

Agarose poly(I)poly(C) precipitation

- 15 To each in vitro kinase reaction mixture, 30 µl of a 50% Ag poly(I)poly(C) Type 6 slurry (Pharmacia LKB Biotechnology) was added, and the mixture was incubated at 4°C for 1 h. The Ag poly(I)poly(C) beads with the absorbed, labelled proteins were then washed four times with was buffer (20 mM HEPES [7.5 pH], 90 mM KCl, 0.1 mM EDTA, 2 mM dithiothreitol, 10% glycerol) at room temperature and mixed with 2X
- 20 Laemmli SDS sample buffer. The beads were then boiled for 5 min, and the released proteins were analyzed by SDS-PAGE.

Polymerase chain reaction

Cells at various times postinfection were harvested and resuspended in ice cold TNE (10 mM Tris [pH 7.8], 150 mM NaCl, 1 mM EDTA) to which NP-40 was then

-14-

added to a final concentration of 1%. After 5 minutes, the nuclei were pelleted and RNA was extracted from the supernatant using the phenol:chloroform procedure. Equal amounts of total cellular RNA from each sample were then subjected to RT-PCR (Wong, H., *et al.*, (1994) *Anal. Biochem.*, 223:251-258) using random hexanucleotide
5 primers (Pharmacia) and RTase (GIBCO-BRL) according to the manufacturers' protocol. The cDNA's from the RT-PCR step was then subjected to selective amplification of reovirus s1 cDNA using the primer 5'-
AATTCGATTAGGTGACACTATAGCTATTGGTCGGATG-3' (SEQ ID NO:1) and
5'-CCCTTTTGACAGTGATGCTCCGTTATCACTCG-3' (SEQ ID NO:2) that amplify
10 a predicted 116 bp fragment. These primer sequences were derived from the S1 sequence determined previously (Nagata, L., *et al.*, (1984) *Nucleic Acids Res.*, 12:8699-8710). The GAPDH primers (Wong, H., *et al.*, (1994) *Anal. Biochem.*, 223:251-258),
5'-CGGAGTCAACGGATTTGGTCGTAT-3' (SEQ ID NO:3) and
5'-AGCCTTCTCCATGGTGGTGAAGAC-3' (SEQ ID NO:4) were used to amplify a
15 predicted 306 bp GAPDH fragment which served as a PCR and gel loading control. Selective amplification of the s1 and GAPDH cDNA's was performed using Taq DNA polymerase (GIBCO-BRL) according to the manufacturers' protocol using a Perkin Elmer Gene Amp PCR system 9600. PCR was carried out for 28 cycles with each
20 consisting of a denaturing step for 30 seconds at 97°C, annealing step for 45 seconds at 55°C, and polymerization step for 60 seconds at 72°C. PCR products were analyzed by electrophoresis through an ethidium bromide-impregnated TAE-2% agarose gel and photographed under ultra-violet illumination with Polaroid 57 film.

Immunoprecipitation and SDS-PAGE analysis

Immunoprecipitation of ³⁵S-labelled reovirus-infected cell lysates with anti-
25 reovirus serotype 3 serum was carried out as previously described (Lee, P.W.K. *et al.*

-15-

(1981) *Virology*, 108:134-146). Immunoprecipitation of ³²P-labelled cell lysates with an anti-PKR antibody (from Dr. Michael Mathews, Cold Spring Harbor) was similarly carried out. Immunoprecipitates were analyzed by discontinuous SDS-PAGE according to the protocol of Laemmli (Laemmli, U.K., (1970) *Nature*, 227:680-685).

5 EXAMPLE 1. Activated Intermediates in the Ras Signalling Pathway Augment
Reovirus Infection Efficiency

It was previously shown that 3T3 cells and their derivatives lacking epidermal growth factor receptors (EGFR) are poorly infectible by reovirus, whereas the same cells transformed with either EGFR or v-erb B are highly susceptible as determined by
10 cytopathic effects, viral protein synthesis, and virus output (Strong, J.E. *et al.*, (1993) *Virology*, 197:405-411; Strong, J.E. and Lee, P.W.K., (1996) *J. Virol.*, 70:612-616).

To determine whether downstream mediators of the EGFR signal transduction pathway may be involved, a number of different NIH 3T3-derived, transformed with constitutively activated oncogenes downstream of the EGFR, were assayed for relative
15 susceptibility to reovirus infection. Of particular interest were intermediates in the ras signalling pathway (reviewed by Barbacid, M., *Annu. Rev. Biochem.*, 56:779-827 (1987); Cahill, M.A., *et al.*, *Curr. Biol.*, 6:16-19 (1996)). To investigate the Ras signalling pathway, NIH 3T3 parental cell lines and NIH 3T3 lines transfected with activated versions of Sos (Aronheim, A., *et al.*, (1994) *Cell*, 78:949-961) or ras
20 (Mundschau, L.J. and Faller, D.V., (1992) *J. Biol. Chem.*, 267:23092-23098) oncogenes were exposed to reovirus, and their capacity to promote viral protein synthesis was compared.

Detection of viral proteins was initially carried out using indirect immunofluorescent microscopy as described above. The results indicated that whereas
25 the NIH 3T3 cells adopted a typically flattened, spread-out morphology with marked

-16-

contact inhibition, the transformed cells all grew as spindle-shaped cells with much less contact inhibition. On comparing the uninfected parental cell lines with the various transformed cell lines, it was apparent that the morphology of the cells was quite distinct upon transformation. Upon challenge with reovirus, it became clear that
5 parental NIH 3T3 line was poorly infectible (<5%), regardless of the source of the parental NIH 3T3 line. In contrast, the transfected cell lines each demonstrated relatively pronounced immunofluorescence by 48 hours postinfection (data not shown).

To demonstrate that viral protein synthesis was more efficient in the Sos- or Ras-transformed cell lines, cells were continuously labeled with [³⁵S]-methionine from
10 12 to 48 hr postinfection and the proteins were analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), as described above.

The results showed clearly that the levels of viral protein synthesis were significantly higher in the Sos- or Ras-transformed cells than in parental NIH 3T3 cells. The identities of the viral bands were confirmed by immunoprecipitation of the labeled
15 proteins with polyclonal anti-reovirus antibodies. Since the uninfected NIH 3T3 cells and their transformed counterparts displayed comparable levels of cellular protein synthesis and doubling times (data not shown), the observed difference in the level of viral protein synthesis could not be due to intrinsic differences in growth rates or translation efficiencies for these cell lines.

20 The long-term fate of infected NIH-3T3 cells was followed by passaging these cells for at least 4 weeks. They grew normally and appeared healthy, with no sign of lytic or persistent infection; no virus could be detected in the medium after this time (data not shown).

-17-

EXAMPLE 2. Enhanced Infectibility Conferred by Activated Oncogenes is Not Due to Long-term Transformation or the Generalized Transformed State of the Cell

To determine whether the differences in susceptibility may be the result of long-term effects of transformation, or the result of the activated oncogene itself, a cell line
5 expressing a zinc-inducible cellular Harvey-ras (c-H-ras) gene was tested for susceptibility to reovirus infectibility, as described above. These cells, called 2H1, were derived from the C3H 10T1/2 cell line which is poorly infectible by reovirus (data not shown), and carry the c-H-ras gene under the control of the mouse metallothionine-I
10 promoter (Trimble, W.S. *et al.* (1986) *Nature*, 321:782-784).

Cells were either mock-treated or pretreated with 50 μ M ZnSO₄ 18 hours prior to infection or mock-infection (administration of carrier agent), followed by indirect immunofluorescent analysis of these cells at 48 hours postinfection or mock-infection.

The results (not shown) demonstrated that uninduced cells were poorly infectible
15 (<5%) whereas those induced for only 18 hours were much more susceptible (>40%). Enhanced viral protein synthesis in the Zn-induced 2H1 cells was further confirmed by metabolic labeling of the cells with [³⁵S]methionine followed by SDS-PAGE analysis of virus-specific proteins (not shown).

Based on these observations, the augmentation of reovirus infection efficiency in
20 the transformed cells is a direct result of the activated oncogene product(s), and not due to other factors such as aneuploidy often associated with long-term transformation, or other accumulated mutations that may be acquired under a chronically transformed state (e.g., p53 or myc activation).

To show further that susceptibility to reovirus infection is not a result of
25 transformation *per se* (i.e., a result of the transformed state of the host cell), NIH-3T3 cells containing a tetracycline-controlled human *c-myc* gene (tet-myc cells) were

-18-

examined (Helbing, C.C. *et al.*, *Cancer Res.* 57:1255-1258 (1997)). These cells normally are maintained in tetracycline (2 µg/ml) which represses the expression of *c-myc*. Removal of tetracycline under normal growth conditions (10% fetal bovine serum) leads to accumulation of the c-Myc protein and the cells display a transformed
5 phenotype. We found that these cells were unable to support virus growth either in the presence or in the absence of tetracycline (data not shown), suggesting that susceptibility to reovirus infection is not due to the general transformed state of the host cell, but rather requires specific transformation by elements of the Ras signaling pathway.

10 A good indicator of activation of the Ras signaling pathway is the activation of the MAP kinases ERK1 and ERK2 (for a review, see Robinson, M.J. and Cobb, M.H., *Curr. Opin. Cell. Biol.* 9:180-186 (1997)). In this regard, it was found that, compared with untransformed cells, Ras-transformed cells have a significantly higher ERK1/2 activity (data not shown). Furthermore, an examination of a number of human cancer
15 cell lines has revealed an excellent correlation between the level of ERK1/2 activity and susceptibility to reovirus infection (data not shown), although ERK1/2 itself does not appear to play any role in it. Mouse L cells and human HeLa cells, in which reovirus grows very well, both manifest high ERK1/2 activity (data not shown).

20 EXAMPLE 3. Viral Transcripts are Generated but Not Translated in Reovirus-Resistant NIH 3T3 Cells

The step at which reovirus infection is blocked in nonsusceptible NIH 3T3 cells was also identified. Because virus binding and virus internalization for nonsusceptible cells were comparable to those observed for susceptible cells (Strong, J.E. *et al.*, (1993) *Virology*, 197:405-411), the transcription of viral genes was investigated.

-19-

The relative amounts of reovirus S1 transcripts generated in NIH 3T3 cells and the Ras-transformed cells during the first 12 hours of infection were compared after amplification of these transcripts by polymerase chain reaction (PCR), as described above. The results demonstrated that the rates of accumulation of S1 transcripts in the two cell lines were similar, at least up to 12 hours postinfection. Similar data were obtained when rates of accumulation of other reovirus transcripts were compared (data not shown). These results demonstrate that infection block in nonsusceptible cells is not at the level of transcription of viral genes, but rather, at the level of translation of the transcripts.

At later times, the level of viral transcripts present in untransformed NIH-3T3 cells decreased significantly whereas transcripts in transformed cells continued to accumulate (data not shown). The inability of these transcripts to be translated in NIH-3T3 cells probably contributed to their degradation. As expected, the level of viral transcripts in infected L cells was at least comparable with that in infected Ras-transformed cells (data not shown).

EXAMPLE 4. A 65 kDa Protein is Phosphorylated in Reovirus-treated NIH 3T3 Cells but Not in Reovirus-infected Transformed Cells

Because viral transcripts were generated, but not translated, in NIH 3T3 cells, it was investigated whether the double-stranded RNA (dsRNA)-activated kinase, PKR, is activated (phosphorylated) in these cells (for example, by S1 mRNA transcripts which have been shown to be potent activators of PKR ((Bischoff, J.R. and Samuel, C.E., (1989) *Virology*, 172:106-115), which in turn leads to inhibition of translation of viral genes. The corollary of such a scenario would be that in the case of the transformed cells, this activation is prevented, allowing viral protein synthesis to ensue.

-20-

NIH 3T3 cells and v-erbB- or Ras- transformed cells (designated THC-11 and H-ras, respectively) were treated with reovirus (i.e., infected) or mock-infected (as above), and at 48 hours post treatment, subjected to *in vitro* kinase reactions, followed by autoradiographic analysis as described above.

- 5 The results clearly demonstrated that there was a distinct phosphoprotein migrating at approximately 65 kDa, the expected size of PKR, only in the NIH 3T3 cells and only after exposure to reovirus. This protein was not labeled in the lysates of either the uninfected transformed cell lines or the infected transformed cell lines. Instead, a protein migrating at approximately 100 kDa was found to be labeled in the transformed
- 10 cell lines after viral infection. This protein was absent in either the preinfection or the postinfection lysates of the NIH 3T3 cell line, and was not a reovirus protein because it did not react with an anti-reovirus serum that precipitated all reovirus proteins (data not shown). A similar 100 kDa protein was also found to be ³²P-labeled in *in vitro* kinase reactions of postinfection lysates of the Sos-transformed cell lines (data not shown).
- 15 That intermediates in the Ras signalling pathway were responsible for the lack of phosphorylation of the 65 kDa protein was further confirmed by the use of the 2H1 cells which contain a Zn-inducible Ras oncogene. Uninduced 2H1 cells (relatively resistant to reovirus infection, as shown above), were capable of producing the 65 kDa phosphoprotein only after exposure to virus. However, 2H1 cells subjected to Zn-
- 20 induction of the H-Ras oncogene showed significant impairment of the production of this phosphoprotein. This impairment coincided with the enhancement of viral synthesis. These results therefore eliminated the possibility that the induction of the 65 kDa phosphoprotein was an NIH 3T3-specific event, and clearly established the role of Ras in preventing (or reversing) induction of the production of this phosphoprotein.
- 25 The Zn-induced 2H1 cells did not produce the 100 kDa phosphoprotein seen in the infected, chronically transformed H-Ras cells.

-21-

EXAMPLE 5. Induction of Phosphorylation of the 65 kDa Protein Requires Active Viral Transcription

Since production of the 65 kDa phosphoprotein occurred only in cells that were resistant to reovirus infection, and only after the cells were exposed to reovirus, it was investigated whether active viral transcription was required for production of the 68 kDa phosphoprotein. Reovirus was UV-treated to inactivate its genome prior to administration of the reovirus to NIH 3T3 cells. For UV-treatment, reovirus was suspended in DMEM to a concentration of approximately 4×10^8 PFU/mL and exposed to short-wave (254 nm) UV light for 20 minutes. UV-inactivated virus were non-infectious as determined by lack of cytopathic effects on mouse L-929 fibroblasts and lack of viral protein synthesis by methods of [^{35}S]-methionine labelling as previously described. Such UV treatment abolished viral gene transcription, as analyzed by PCR, and hence viral infectivity (data not shown). The cells were then incubated for 48 hours, and lysates were prepared and subjected to *in vitro* ^{32}P -labeling as before. The results showed that NIH 3T3 cells infected with untreated reovirus produced a prominent 65 kDa ^{32}P -labelled band not found in uninfected cells. Cells exposed to UV-treated reovirus behaved similarly to the uninfected control cells, manifesting little phosphorylation of the 65 kDa protein. Thus, induction of the phosphorylation of the 65 kDa phosphoprotein is not due to dsRNA present in the input reovirus; rather, it requires *de novo* transcription of the viral genes, consistent with the identification of the 65 kDa phosphoprotein as PKR.

EXAMPLE 6. Identification of the 65 kDa Phosphoprotein as PKR

To determine whether the 65 kDa phosphoprotein was PKR, a dsRNA binding experiment was carried out in which poly(I)-poly(c) agarose beads were added to ^{32}P -labeled lysates, as described above. After incubation for 30 minutes at 4°C , the beads

-22-

were washed, and bound proteins were released and analyzed by SDS-PAGE. The results showed that the 65 kDa phosphoprotein produced in the postinfection NIH 3T3 cell lysates was capable of binding to dsRNA; such binding is a well-recognized characteristic of PKR. In contrast, the 100 kDa phosphoprotein detected in the infected H-ras-transformed cell line did not bind to the Poly(I)-poly(c) agarose. The 65 kDa phosphoprotein was also immunoprecipitable with a PKR-specific antibody (provided by Dr. Mike Mathews, Cold Spring Harbor Laboratory), confirming that it was indeed PKR.

EXAMPLE 7. PKR Inactivation or Deletion Results in Enhanced Infectibility of Untransformed Cells

If PKR phosphorylation is responsible for the shut-off of viral gene translation in NIH-3T3 cells, and one of the functions of the activated oncogene product(s) in the transformed cells is the prevention of this phosphorylation event, then inhibition of PKR phosphorylation in NIH-3T3 cells by other means (e.g. drugs) should result in the enhancement of viral protein synthesis, and hence infection, in these cells. To test this idea, 2-aminopurine was used. This drug has been shown to possess relatively specific inhibitory activity towards PKR autophosphorylation (Samuel, C.E. and Brody, M., (1990) *Virology*, 176:106-113; Hu, Y. and Conway, T.W. (1993), *J. Interferon Res.*, 13:323-328). Accordingly, NIH 3T3 cells were exposed to 5 mM 2-aminopurine concurrently with exposure to reovirus. The cells were labeled with [³⁵S]methionine from 12 to 48 h postinfection, and lysates were harvested and analyzed by SDS-PAGE.

The results demonstrated that exposure to 2-aminopurine resulted in a significantly higher level of viral protein synthesis in NIH 3T3 cells (not shown). The enhancement was particularly pronounced after immunoprecipitating the lysates with an anti-reovirus serum. These results demonstrate that PKR phosphorylation leads to

-23-

inhibition of viral gene translation, and that inhibition of this phosphorylation event releases the translation block. Therefore, intermediates in the Ras signalling pathway negatively regulate PKR, leading to enhanced infectibility of Ras-transformed cells.

Interferon β , known to induce PKR expression, was found to significantly
5 reduce reovirus replication in Ras-transformed cells (data not shown).

A more direct approach to defining the role of PKR in reovirus infection is through the use of cells that are devoid of PKR. Accordingly, primary embryo fibroblasts from wild-type PKR $^{+/+}$ and PKR $^{-/-}$ mice (Yang, Y.L. *et al. EMBO J.* 14:6095-6106 (1995)) were compared in terms of susceptibility to reovirus infection.
10 The results clearly showed that reovirus proteins were synthesized at a significantly higher level in the PKR $^{-/-}$ cells than in the PKR $^{+/+}$ cells. These experiments demonstrated that PKR inactivation or deletion enhanced host cell susceptibility to reovirus infection in the same way as does transformation by Ras or elements of the Ras signaling pathway, thereby providing strong support of the role of elements of the Ras
15 signaling pathway in negatively regulating PKR.

EXAMPLE 8 Inactivation of PKR in Transformed Cells Does Not Involve MEK

Receptor tyrosine kinases such as EGFRs are known to stimulate the mitogen-activated or extracellular signal-regulated kinases (ERK1/2) via Ras (see Robinson, M.J. and Cobb, M.H., *Curr. Opin. Cell. Biol.* 9:180-186 (1997)). This stimulation
20 requires the phosphorylation of ERK1/2 by the mitogen-activated extracellular signal-regulated kinase, kinase MEK, which itself is activated (phosphorylated) by Raf, a serine-threonine kinase downstream of Ras. To determine if MEK activity was required for PKR inactivation in transformed cells, the effect of the recently identified EK inhibitor PD98059 (Dudley, D.T. *et al., Proc. Natl. Acad. Sci. USA* 92:7686-7689

-24-

(1995); Waters, S.D. *et al.*, *J. Biol. Chem.* 270:20883-20886 (1995)) on infected Ras-transformed cells was studied.

H-Ras-transformed cells were grown to 80% confluency and infected with reovirus at an m.o.i. of approximately 10 p.f.u./cell. PD98059 (Calbiochem), dissolved
5 in dimethylsulfoxide (DMSO), was applied to the cells at the same time as the virus (final concentration of PD98059 was 50 μ M). The control cells received an equivalent volume of DMSO. The cells were labeled with ³⁵S-methionine from 12 to 48 hours post-infection. Lysates were then prepared, immunoprecipitated with the polyclonal anti-reovirus serotype 3 serum and analyzed by SDS-PAGE.

10 The results (data not shown) showed that PD98059, at a concentration that effectively inhibited ERK1/2 phosphorylation, did not inhibit reovirus protein synthesis in the transformed cells. On the contrary, PD98059 treatment consistently caused a slight enhancement of viral protein synthesis in these cells; the reason for this is under investigation. Consistent with the lack of inhibition of viral protein synthesis in the
15 presence of PD98059, the PKR in these cells remained unphosphorylated (data not shown). As expected, PD98059 had no effect on reovirus-induced PKR phosphorylation in untransformed NIH-3T3 cells (data not shown). These results indicated that MEK and ERK1/2 are not involved in PKR activation.

EXAMPLE 9. In Vivo Oncolytic Capability of Reovirus

20 A severe combined immunodeficiency (SCID) host tumor model was used to assess the efficacy of utilizing reovirus for tumor reduction. Male and female SCID mice (Charles River, Canada) were injected with v-erbB-transformed NIH 3T3 mouse fibroblasts (designated THC-11 cells) in two subcutaneous sites overlying the hind flanks. In a first trial, an injection bolus of 2.3×10^5 cells in 100 μ l of sterile PBS was

-25-

used. In a second trial, an injection bolus of 4.8×10^6 cells in 100 μ l PBS was used. Palpable tumors were evident approximately two to three weeks post injection.

Reovirus serotype three (strain Dearing) was injected into the right-side tumor mass (the "treated tumor mass") in a volume of 20 μ l at a concentration of 1.0×10^9 plaque forming units (PFU)/ml. The left-side tumor mass (the "untreated tumor mass") was left untreated. The mice were observed for a period of seven days following injection with reovirus, measurements of tumor size were taken every two days using calipers, and weight of tumors was measured after sacrifice of the animals. All mice were sacrificed on the seventh day. Results are shown in Table 1.

10 Table 1 Tumor Mass after Treatment with Reovirus

Trial 1 (n=8)	mean untreated tumor mass	602 mg
	mean treated tumor mass	284 mg
Trial 2 (n = 12)		
	mean control tumor mass	1523.5 mg
	mean untreated tumor mass	720.9 mg
	mean treated tumor mass	228.0 mg

The treated tumor mass was 47% of that of the untreated tumor mass in trial 1, and 31.6% of the untreated tumor mass in trial 2. These results indicated that the virus-treated tumors were substantially smaller than the untreated tumors, and that there may be an additional systemic effect of the virus on the untreated tumor mass.

Similar experiments were also conducted using unilateral introduction of tumor cells. SCID mice were injected subcutaneously and unilaterally in the hind flank with v-erbB-transformed NIH 3T3 mouse fibroblasts (THC-11 cells). Palpable tumors (mean area 0.31 cm²) were established after two weeks. Eight animals were then given

-26-

a single intratumoral injection of 1.0×10^7 PFUs of reovirus serotype 3 (strain Dearing) in phosphate-buffered saline (PBD). Control tumors (n=10) were injected with equivalent amounts of UV-inactivated virus. Tumor growth was followed for 12 days, during which time no additional reovirus treatment was administered.

- 5 Results, shown in Figure 2, demonstrated that treatment of these tumors with a single dose of active reovirus (open circles) resulted in dramatic repression of tumor growth by the thirteenth day (endpoint), when tumors in the control animals injected with a single dose of inactivated reovirus (closed circles) exceeded the acceptable tumor burden. This experiment was repeated several times and found to be highly
10 reproducible, thus further demonstrating the efficacy of reovirus in repressing tumor growth.

Example 10: *In Vivo* Oncolytic Capability of Reovirus Against Human Breast Cancer-Derived Cell Lines

- In vivo* studies were also carried out using human breast carcinoma cells in a
15 SCID mouse model. Female SCID mice were injected with 1×10^9 MDAMB468 cells in two subcutaneous sites, overlying both hind flanks. Palpable tumors were evident approximately two to four weeks post injection. Undiluted reovirus serotype three (strain Dearing) was injected into the right side tumor mass in a volume of 20 μ l at a concentration of 1.0×10^{10} PFU/ml. The results are shown in Table 2.

-27-

Table 2 Tumor Mass After Treatment with Reovirus

TREATMENT	mean untreated tumor mass (left side)	mean treated tumor mass (right side)
Reovirus (N=8)	29.02 g	38.33 g
Control (N=8)	171.8 g	128.54 g

- 5 *Note: One of the control mice died early during the treatment phase. None of the reovirus-treated mice died.

Although these studies were preliminary, it was clear that the size of the tumors in the reovirus-treated animals was substantially lower than that in the untreated animals. However, the size of the tumors on the right (treated) side of the reovirus-
10 treated animals was slightly larger on average than the left (untreated) side. This was unexpected but may be explained by the composition of the mass being taken up by inflammatory cells with subsequent fibrosis, as well as by the fact that these tumors were originally larger on the right side on average than the left. The histologic composition of the tumor masses is being investigated. These results also support the
15 systemic effect the reovirus has on the size of the untreated tumor on the contralateral slide of reovirus injection.

EXAMPLE 11. Susceptibility of Additional Human Tumors to Reovirus Oncolysis

In view of the *in vivo* results presented above, the oncolytic capability observed
20 in murine cells was investigated in cell lines derived from additional human tumors.

A. Materials and Methods

-28-

Cells and Virus

All cell lines were grown in Dulbecco's modified Eagle's medium (DMEM) containing 10% fetal bovine serum (FBS).

The Dearing strain of reovirus serotype 3 used in these studies was propagated
5 in suspension cultures of L cells and purified according to Smith (Smith, R.E. *et al.*,
(1969) *Virology*, 39:791-800) with the exception that β -mercaptoethanol (β -ME) was
omitted from the extraction buffer. Reovirus labelled with [35 S]methionine was grown
and purified as described by McRae and Joklik (McRae, M.A. and Joklik, W.K., (1978)
Virology, 89:578-593). The particle/PFU ration for purified reovirus was typically
10 100/1.

Cytopathic effects of reovirus on cells

Confluent monolayers of cells were infected with reovirus serotype 3 (strain
Dearing) at a multiplicity of infection (MOI) of approximately 40 plaque forming units
(PFU) per cell. Pictures were taken at 36 hour postinfection for both reovirus-infected
15 and mock-infected cells.

Immunofluorescent analysis of reovirus infection

For the immunofluorescent studies the cells were grown on coverslips, and
infected with reovirus at a multiplicity of infection (MOI) of ~ 10 PFU/cell or mock-
infected as described above. At various times postinfection, cells were fixed in an
20 ethanol/acetic acid (20/1) mixture for 5 minutes, then rehydrated by subsequential
washes in 75%, 50% and 25% ethanol, followed by 4 washes with phosphate-buffered
saline (PBS). The fixed and rehydrated cells were then exposed to the primary antibody
(rabbit polyclonal anti-reovirus type 3 serum diluted 1/100 in PBS) for 2 hr at room
temperature. Following 3 washes with PBS, the cells were exposed to the secondary

-29-

antibody [goat anti-rabbit IgG (whole molecule) fluorescein isothiocyanate (FITC) conjugate diluted 1/100 in PBS containing 10% goat serum and 0.005% Evan's Blue counterstain] for 1 hour at room temperature. Finally, the fixed and treated cells were washed 3 more times with PBS, followed by 1 wash with double-distilled water, dried
5 and mounted on slides in 90% glycerol containing 0.1% phenylenediamine, and viewed with a Zeiss Axiophot microscope mounted with a Carl Zeiss camera (magnification for all pictures was 200 x).

Infection of cells and quantitation of virus

Confluent monolayers of cells grown in 24-well plates were infected with
10 reovirus at an estimated multiplicity of 10 PFU/cell. After 1 hour incubation at 37°C, the monolayers were washed with warm DMEM-10% FBS, and then incubated in the same medium. At various times postinfection, a mixture of NP-40 and sodium deoxycholate was added directly to the medium on the infected monolayers to final concentrations of 1% and 0.5%, respectively. The lysates were then harvested and virus
15 yields were determined by plaque titration on L-929 cells.

Radiolabelling of reovirus-Infected cells and preparation of lysates

Confluent monolayers of cells were infected with reovirus (MOI ~10 PFU/cell). At various times postinfection, the media was replaced with methionine-free DMEM containing 10% dialyzed PBS and 0.1 mCi/ml [³⁵S]methionine. After further incubation
20 for 1 hour at 37°C, the cells were washed in phosphate-buffered saline (PBS) and lysed in the same buffer containing 1% Triton X-100, 0.5% sodium deoxycholate and 1 mM EDTA. The nuclei were then removed by low speed centrifugation and the supernatants was stored at 70°C until use.

-30-

Immunoprecipitation and SDS-PAGE analysis

Immunoprecipitation of [³⁵S]-labelled reovirus-infected cell lysates with anti-reovirus serotype 3 serum was carried out as previously described (Lee, P.W.K. *et al.* (1981) *Virology*, 108:134-146). Immunoprecipitates were analyzed by discontinuous
5 SDS-PAGE according to the protocol of Laemmli (Laemmli, U.K., (1970) *Nature*, 227:680-685).

B. Breast Cancer

The *c-erbB-2/neu* gene encodes a transmembrane protein with extensive homology to the EGFR that is overexpressed in 20-30% of patients with breast cancer
10 (Yu, D. *et al.* (1996) *Oncogene* 13:1359). Since it has been established herein that Ras activation, either through point mutations or through augmented signaling cascade elements upstream of Ras (including the *c-erbB-2/neu* homologue EGFR) ultimately creates a hospitable environment for reovirus replication, an array of cell lines derived from human breast cancers were assayed for reovirus susceptibility. The cell lines
15 included MDA-MD-435SD (ATCC deposit HTB-129), MCF-7 (ATCC deposit HTB-22), T-27-D (ATCC deposit HTB-133), BT-20 (ATCC deposit HTB-19), HBL-100 (ATCC deposit HTB-124), MDA-MB-468 (ATCC deposit HTB-132), and SKBR-3 (ATCC deposit HTB-30).

Based upon induction of cytopathic effects, and viral protein synthesis as
20 measured by radioactive metabolic labeling and immunofluorescence as described above, it was found that five out of seven of the tested breast cancers were susceptible to reovirus infection: MDA-MB-435S, MCF-7, T-27-D, MDA MB-468, and SKBR-3 were exquisitely sensitive to infection, while BT-20 and HBL-100 demonstrated no infectibility.

-31-

C. Brain Glioblastoma

Next a variety of cell lines derived from human brain glioblastomas was investigated. The cell lines included A-172, U-118, U-178, U-563, U-251, U-87 and U-373 (cells were a generous gift from Dr. Wee Yong, University of Calgary).

5 Six out of seven glioblastoma cell lines demonstrated susceptibility to reovirus infection, including U-118, U-178, U-563, U-251, U-87 and U-373, while A-172 did not demonstrate any infectibility, as measured by cytopathic effects, immunofluorescence and [³⁵S]-methionine labeling of reovirus proteins.

10 The U-87 glioblastoma cell line was investigated further. To assess the sensitivity of U-87 cells to reovirus, U-87 cells (obtained from Dr. Wee Yong, University of Calgary) were grown to 80% confluency and were then challenged with reovirus at a multiplicity of infection (MOI) of 10. Within a period of 48 hours there was a dramatic, widespread cytopathic effect (data not shown). To demonstrate further
15 that the lysis of these cells was due to replication of reovirus, the cells were then pulse-labeled with [³⁵S]methionine for three hour periods at various times postinfection and proteins were analyzed by sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE) as described above. The results (not shown) clearly demonstrated effective reovirus replication within these cells with resultant shutoff of host protein
20 synthesis by 24 hours postinfection.

The U-87 cells were also introduced as human tumor xenografts into the hind flank of 10 SCID mice. U-87 cells were grown in Dulbecco's modified Eagle's medium containing 10% fetal bovine serum, as described above. Cells were harvested, washed, and resuspended in sterile PBS; 2.0×10^6 cells in 100 μ l were injected
25 subcutaneously at a site overlying the hind flank in five- to eight-week old male SCID mice (Charles River, Canada). Tumor growth was measured twice weekly for a period

-32-

of four weeks. Results, shown in Figure 3, demonstrated that treatment of U-87 tumors with a single intratumoral injection of 1.0×10^7 PFUs of live reovirus (open circles, $n=5$) resulted in drastic repression of tumor growth, including tumor regression by the fourth week post-treatment ($P=0.008$), in comparison with treatment of tumors with a
5 single intratumoral injection of the same amount of UV-inactivated reovirus (closed circles, $n=5$).

Hematoxylin/eosin (HE)-staining of the remaining microfoci of the tumors treated with active virus, performed as described (H. Lyon, *Cell Biology, A Laboratory Handbook*, J.E. Celis, ed., Academic Press, 1994, p. 232) revealed that the remaining
10 tumor mass consisted largely of normal stroma without appreciable numbers of viable tumor cells, nor was there any evidence of infiltration of tumor cells into the underlying skeletal muscle (data not shown). Necrosis of tumor cells was due to direct lysis by the virus, the same mechanism of cell killing as by reovirus *in vitro*.

To determine if there was viral spread beyond the tumor mass,
15 immunofluorescent microscopy using antibodies directed against total reovirus proteins was conducted as described above, on sections of the tumor and adjoining tissue. It was found that reovirus-specific proteins were confined to the tumor mass; no viral staining was detected in the underlying skeletal muscle (data not shown). As expected, viral proteins were not present in tumors injected with the UV-inactivated virus (data not
20 shown). These results demonstrated that reovirus replication in these animals was highly tumor specific with viral amplification only in the target U-87 cells.

Since most tumors are highly vascularized, it was likely that some virus could enter the blood stream following the lysis of the infected tumor cells. To determine if there was systemic spread of the virus, blood was harvested from the treated and control
25 animals, and serially diluted for subsequent plaque titration. Infectious virus was found to be present in the blood at a concentration of 1×10^5 PFUs/ml (data not shown).

-33-

The high degree of tumor specificity of the virus, combined with systemic spread, suggested that reovirus could be able to replicate in glioblastoma tumors remote from the initially infected tumor, as demonstrated above with regard to breast cancer cells. To verify this hypothesis, SCID mice were implanted bilaterally with U-87
5 human tumor xenografts on sites overlying each hind flank of the animals. These tumors were allowed to grow until they measured 0.5 x 0.5 cm. The left-side tumors were then administered a single infection of reovirus in treated animals (n=5); control animals (n=7) were mock-treated with UV-inactivated virus. Tumors were again measured twice weekly for a period of four weeks.

10 Results, shown in Figure 4, demonstrated that inhibition and eventual regression of both the treated (circles) and untreated tumor masses (squares) occurred only in the live reovirus-treated animals (open circles and squares), in contrast with the inactivated reovirus-treated animals (closed circles and squares). Subsequent immunofluorescent analysis revealed that reovirus proteins were present in both the ipsilateral (treated) as
15 well as the contralateral (untreated) tumor, indicating that regression on the untreated side was a result of reovirus oncolysis (data not shown).

D. Pancreatic Carcinoma

Cell lines derived from pancreatic cancer were investigated for their susceptibility to reovirus infection. The cell lines included Capan-1 (ATCC deposit
20 HTB-79), BxPC3 (ATCC deposit CRL-1687), MIAPACA-2 (ATCC deposit CRL-1420), PANC-1 (ATCC deposit CRL-1469), AsPC-1 (ATCC deposit CRL-1682) and Hs766T (ATCC deposit HTB-134).

Five of these six cell lines demonstrated susceptibility to reovirus infection including Capan-1, MIAPACA-2, PANC-1, AsPC-1 and Hs766T, whereas BxPC3
25 demonstrated little infectability as assayed by virus-induced cytopathological effects,

-34-

immunofluorescence and [³⁵S]-labelling. Interestingly, four of the five cell lines demonstrating susceptibility to reovirus oncolysis have been shown to possess transforming mutations in codon 12 of the K-ras gene (Capan-1, MIAPACA-2, PANC-1 and AsPC-1) whereas the one lacking such susceptibility (BxPC3) has been shown to
5 lack such a mutation (Berrozpe, G., *et al.* (1994), *Int. J. Cancer*, 58:185-191). The status of the other K-ras codons is currently unknown for the Hs766T cell line.

EXAMPLE 12. Use of Reovirus as an Oncolytic Agent in Immune-Competent
Animals

A syngeneic mouse model was developed to investigate use of reovirus in
10 immune-competent animals rather than in SCID mice as described above. C3H mice (Charles River) were implanted subcutaneously with 1.0×10^7 PFUs ras-transformed C3H cells (a gift of D. Edwards, University of Calgary). Following tumor establishment, mice were treated with a series of injections of either live reovirus (1.0×10^8 PFUs) or UV-inactivated reovirus. Following an initial series (six injections over a
15 nine-day course), test animals received a treatment of dilute reovirus (1.0×10^7 PFUs) every second day. Mock-treated animals received an equivalent amount of UV-inactivated virus.

Results demonstrated that reovirus was an effective oncolytic agent in these immune competent animals. All of the test animals showed regression of tumors; 5 of
20 the 9 test animals exhibited complete tumor regression after 22 days, a point at which the control animals exceeded acceptable tumor burden. Furthermore, there were no identifiable side effects in the animals treated with reovirus.

To assess the effects of previous reovirus exposure on tumor repression and regression, one-half of a test group was challenged with reovirus (intramuscular
25 injection of 1.0×10^8 PFUs, type 3 Dearing) prior to tumor establishment. Two weeks

-35-

after challenge, neutralizing antibodies could be detected in all exposed animals.

Following tumor establishment, animals were treated with a series of either live or UV-inactivated reovirus, as described above.

Results (data not shown) demonstrated that animals with circulating neutralizing
5 antibodies to reovirus (i.e., those challenged with reovirus prior to tumor establishment) exhibited tumor repression and regression similar to those animals in which there was no prior exposure to reovirus. Thus, reovirus can serve as an effective oncolytic agent even in immune-competent animals with previous exposure to reovirus.

While this invention has been particularly shown and described with references
10 to preferred embodiments thereof, it will be understood by those skilled in the art that various changes in form and details may be made therein without departing from the spirit and scope of the invention as defined by the appended claims.

-36-

CLAIMS

What is claimed is:

1. Use of a reovirus for the manufacture of a medicament for treating a Ras-mediated neoplasm in a mammal.
- 5 2. The use of Claim 1, wherein the reovirus is a human reovirus.
3. The use of Claim 2, wherein the human reovirus is selected from the group consisting of: type 1 reovirus, type 2 reovirus, and type 3 reovirus.
4. The use of Claim 1, wherein the reovirus is a nonhuman reovirus.
- 10 5. The use of Claim 4, wherein the nonhuman reovirus is selected from the group consisting of: mammalian reovirus and avian reovirus.
6. The use of Claim 1, wherein more than one type of reovirus is administered.
7. The use of Claim 1, wherein more than one strain of reovirus is administered.
8. The use of Claim 1, wherein the reovirus is a field isolate..
- 15 9. The use of Claim 1, wherein the reovirus is treated with a protease prior to administration.

-37-

10. The use of Claim 1, wherein the neoplasm is a solid neoplasm.
11. The use of Claim 1, wherein the neoplasm is a hematopoietic neoplasm.
12. The use of Claim 1, wherein the mammal is selected from the group consisting of: mice, dogs, cats, sheep, goats, cows, horses, pigs, and non-human primates.
- 5 13. The use of Claim 1, wherein the mammal is a human.
14. The use of Claim 1, wherein the neoplasm is selected from the group consisting of: pancreatic cancer, breast cancer and brain cancer.
15. The use of Claim 1, wherein the neoplasm is selected from the group consisting of: lung cancer, prostate cancer, colorectal cancer, thyroid cancer, renal cancer,
10 adrenal cancer, liver cancer, and leukemia.
16. The use of Claim 1, wherein the ras-mediated neoplasm is metastatic.
17. A method of treating a Ras-mediated neoplasm in a mammal, comprising administering to the neoplasm a reovirus in an amount sufficient to result in reovirus-mediated oncolysis of cells of the neoplasm.
- 15 18. The method of Claim 17, wherein the reovirus is a human reovirus.
19. The method of Claim 18, wherein the human reovirus is selected from the group consisting of: type 1 reovirus, type 2 reovirus, and type 3 reovirus.

-38-

20. The method of Claim 16, wherein the reovirus is a non-human reovirus.
21. The method of Claim 20, wherein the reovirus is selected from the group consisting of: mammalian reovirus and avian reovirus.
22. The method of Claim 17, wherein more than one type of reovirus is
5 administered.
23. The method of Claim 17, wherein more than one strain of reovirus is administered.
24. The method of Claim 17, wherein the reovirus is a field isolate.
25. The method of Claim 17, wherein the reovirus is treated with a protease prior to
10 administration.
26. The method of Claim 17, wherein the neoplasm is a solid neoplasm.
27. The method of Claim 26, wherein the reovirus is administered by injection into the solid neoplasm.
28. The method of Claim 26, wherein the reovirus is administered intravenously into
15 the mammal.
29. The method of Claim 17, wherein the neoplasm is a hematopoietic neoplasm.

-39-

30. The method of Claim 29, wherein the reovirus is administered intravenously into the mammal.
31. The method of Claim 29, wherein the reovirus is administered intraperitoneally into the mammal.
- 5 32. The method of Claim 17, wherein the mammal is selected from the group consisting of: mice, dogs, cats, sheep, goats, cows, horses, pigs, and non-human primates.
33. The method of Claim 17, wherein the mammal is a human.
34. The method of Claim 17, wherein the neoplasm is selected from the group
10 consisting of: pancreatic cancer, breast cancer and brain cancer.
35. The method of Claim 17, wherein the neoplasm is selected from the group consisting of: lung cancer, prostate cancer, colorectal cancer, thyroid cancer, renal cancer, adrenal cancer, liver cancer, and leukemia.
36. The method of Claim 17, wherein approximately 10^3 to 10^{12} plaque forming
15 units of reovirus are administered.
37. The method of Claim 17, wherein the reovirus is administered in a single dose.
38. The method of Claim 17, wherein the reovirus is administered in more than one dose.

-40-

39. The method of Claim 17, wherein the reovirus is administered to more than one neoplasm in the mammal.
40. The method of Claim 17, wherein the ras-mediated neoplasm is metastatic.
41. The method of Claim 40, wherein the reovirus is administered to a single solid
5 neoplasm.
42. The method of Claim 40, wherein the reovirus is administered intravenously.
43. A method of treating a Ras-mediated neoplasm in a human, comprising administering to the neoplasm a reovirus in an amount sufficient to result in reovirus-mediated oncolysis of cells of the neoplasm.

1/4




Figure 1

2/4




Figure 2

3/4




Figure 3

4/4




Figure 4

INTERNATIONAL SEARCH REPORT

In national Application No

PCT/CA 98/00774

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 A61K35/76

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	G. HASHIRO ET AL.: "THE PREFERENTIAL CYTOTOXICITY OF REOVIRUS FOR CERTAIN TRANSFORMED CELL LINES." ARCHIVES OF VIROLOGY, vol. 54, no. 4, 1977, pages 307-315, XP002084838 VIENNA, AT see the whole document ----- -/--	1-43



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier document but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

"&" document member of the same patent family

Date of the actual completion of the international search

18 November 1998

Date of mailing of the international search report

02/12/1998

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Ryckebosch, A

INTERNATIONAL SEARCH REPORT

International Application No

PCT/CA 98/00774

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	J.E. STRONG ET AL.: "THE v-erbB ONCOGENE CONFERS ENHANCED CELLULAR SUSCEPTIBILITY TO REOVIRUS INFECTION." JOURNAL OF VIROLOGY., vol. 70, no. 1, January 1996, pages 612-616, XP002084839 WASHINGTON, DC, US cited in the application see page 615, right-hand column, line 28 - page 616, left-hand column, line 6 ----	1-43
A	WO 94 18992 A (ONYX PHARMACEUTICALS) 1 September 1994 see claims ----	1-43
A	EP 0 514 603 A (THE GENERAL HOSPITAL CORPORATION) 25 November 1992 see claims ----	1-43
P,X	J.E. STRONG ET AL.: "THE MOLECULAR BASIS OF VIRAL ONCOLYSIS: USURPATION OF THE Ras SIGNALING PATHWAY BY REOVIRUS." THE EMBO JOURNAL., vol. 17, no. 12, 15 June 1998, pages 3351-3362, XP002084840 EYNSHAM, OXFORD GB see the whole document -----	1-43

INTERNATIONAL SEARCH REPORT

International application No.

PCT/CA 98/ 00774

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Remark: Although claims 17-43 are directed to a method of treatment of the human/animal body, the search has been carried out and based on the alleged effect of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

☐ The additional search fees were accompanied by the applicant's protest.

☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

In ternational Application No

PCT/CA 98/00774

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9418992 A	01-09-1994	AU 682854 B	23-10-1997
		AU 6272294 A	14-09-1994
		CA 2152941 A	01-09-1994
		EP 0689447 A	03-01-1996
		JP 8507076 T	30-07-1996
		US 5801029 A	01-09-1998
		US 5677178 A	14-10-1997
EP 514603 A	25-11-1992	CA 2051289 A	15-03-1992
		JP 6100458 A	12-04-1994